

**EFFECT OF EFFECTIVE MICROORGANISM (EM) ON GROWTH AND
YIELD OF AEROBIC RICE VARIETY MR1A1**

MUHAMMAD HAFIZ BIN HASAN

**Final Year Project Report Submitted in
Partial Fulfilment of the Requirement for the
Degree of Bachelor of Science (Hons.) Plantation Management and Technology
in the Faculty of Plantation and Agrotechnology
Universiti Teknologi MARA**

JULY 2015

DECLARATION

This Final Year Project is a partial fulfillment of the requirements for a degree of Bachelor of Science (Hons.) Technology and Plantation Management, Faculty of Plantation and Agrotechnology, Universiti Teknologi MARA.

It is entirely my own work and has not been submitted to any other University or higher education institution, or for any other academic award in this University. Where use has been made of the work of other people it has been fully acknowledged and fully referenced.

I hereby assign all and every right in the copyright to this Work to the Universiti Teknologi MARA ("UiTM"), which henceforth shall be the owner of copyright in this Work and that, any reproduction or use in any form or by any means whatsoever is prohibited without a written consent of UiTM.

Candidate's signature: Date:

Name: MUHAMMAD HAFIZ BIN HASAN

I hereby declare that I have checked this project and in my opinion, this project is adequate in terms of scope and quality for the award of the degree of Bachelor of Science (Hons.) Technology and Plantation Management, Faculty of Plantation and Agrotechnology, Universiti Teknologi MARA.

Signature:

Name of Supervisor: PROF MADYA DR ADZMI BIN YAACOB

Position: SENIOR LECTURER

Date:

TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGEMENT	iii
TABLE OF CONTENTS	iv
LIST OF FIGURES	vi
LIST OF TABLES	vii
LIST OF ABBREVIATIONS	viii
ABSTRACT	ix
ABSTRAK	x
<u>CHAPTER</u>	
1 INTRODUCTION	
1.1 Background of study	1
1.2 Problem statement	2
1.3 Significance of the Study	2
1.4 Objective of the Study	3
2 LITERATURE REVIEW	
2.1 Aerobic rice	4
2.2 Variety of aerobic rice (MRJA1)	4
2.3 Effective microorganism (EM)	5
2.4 Soil chemical properties	
2.4.1 Nutrient contents	5
2.4.2 pH value	6
2.5 Microbial population	6
3 METHODOLOGY	
3.1 Location of study	7
3.2 Material and equipment	
3.2.1 Effective microorganism (EM-1)	7
3.3.2 Rice variety	7
3.3.3 Apparatus	7
3.3.4 Pots	7
3.3 Parameter	
3.3.1 Plant height measurement	8
3.3.2 Number of tiller	8
3.3.3 Number of leaves	8
3.3.4 Number of panicle	8
3.3.5 Soil content	8
3.3.6 Soil pH	8
3.3.7 Population of microbial (bacteria)	8
3.4 Experimental Procedures	
3.4.1 Preparation of rice seedling	9
3.4.2 Preparation of effective microorganism	9
3.4.3 Preparation of treatment	10
3.4.4 Determination of soil nutrient content	12

	3.4.5	Determination of soil pH	13
	3.4.6	Serial dilution (different rate of EM concentration and applied soil)	13
	3.4.7	Microbial population counting	14
	3.5	Experimental design	15
	3.6	Analysis of statistical	15
	3.7	Schedule of framework	16
4	RESULTS		
	4.1	Effect of different rate of EM concentration on plant height	18
	4.2	Effect of different rate of EM concentration on number of tillers	20
	4.3	Effect of different rate of EM concentration on number of leaves	22
	4.4	Effect of different rate of EM concentration on number of panicle	24
	4.5	Effect of different rate of EM concentration on soil nutrient contents	25
	4.6	Effect of different rate of EM concentration on soil pH value	27
	4.7	Population of bacteria in different concentration of EM	29
	4.8	Population of bacteria in soil	30
5	DISCUSSION		
	5.1	Discussion	31
6	CONCLUSION AND RECOMMENDATION		
	5.1	Conclusion	33
	5.2	Recommendation	33
	REFERENCES		35
	APPENDICES		37
	CURRICULUM VITAE		51

ABSTRACT

EFFECT OF EFFECTIVE MICROORGANISM (EM) ON GROWTH AND YIELD OF AEROBIC RICE VARIETY MR1A1

Rice is the staple food for Malaysia. Rice production in the country is insufficient to meet the demand of our population. To overcome the shortage Malaysia had to import rice from country like Thailand. An abandoned land commonly found in the country can be utilized for aerobic rice cultivation. Other than that, to overcome the fertility problem of abandoned lands, inorganic fertilizers can be used together with EM. In this study, the effective microorganisms improve soil chemical properties, soil fertility and plant where they do activities of microorganisms in the soil. EM that originating from liquid in which is consist of various species of microorganisms such as lactic acid bacteria, photosynthetic bacteria, yeast, and others. The study was conducted in a greenhouse at UiTM Jasin by making 5 treatments T1, T2, T3, T4 and T5 (T1 = NPK, T2 = EM 1:500, T3 = EM 1:700, T4 = EM 1:900 and T5 = NPK + EM 1:300). The observation of this study was conducted for 9 weeks until the plant reaches the stage of reproductive (70 – 90 days). Besides that, the identification and estimation of the bacterial population in liquid concentration of effective microorganism (EM) was identified by differences in CFU in each treatment of EM and EM in the soil. The results show that NPK fertilizers is more effective for growth of rice than EM. All EM treatment showed no significant difference between all of them in terms of plant height, number of tiller, the number of leaves, and number of panicle. However, there is significant difference in nutrient content and pH value in the soil. Although EM is less effective than NPK fertilizers in terms of overall growth, they seem to improve the activities of microorganisms in the soil. In the long run EM has a potential to improve the yield of rice.